

Higher-dimensional operators in SUSY SO(10) GUT models

J. Sayre and S. Wiesenfeldt

Department of Physics, University of Illinois at Urbana-Champaign,
1110 West Green Street, Urbana, IL 61801, USA

Abstract

SO(10) GUT models with only small Higgs fields use higher-dimensional operators to generate realistic fermion mass matrices. In particular, a Higgs field in the spinor representation, 16_H^d , acquires a weak scale vev. We include the weak vev of the corresponding field $\bar{16}_H^u$ and investigate its effects based on two successful models, one by Albright and Barr (AB) and another by Babu, Pati and Wilczek (BPW). We find that the BPW model is a particular case within a class of models with identical fermion masses and mixings. In contrast, we expect corrections to the parameters of AB-type models.

Introduction. SO(10) is the most attractive candidate for a Grand Unified Theory (GUT): The matter particles of each standard model generation, together with one singlet, the right-handed neutrino, are unified in a single representation of a simple gauge group. The breaking of supersymmetric SO(10) at $M_{\text{GUT}} \simeq 2 \cdot 10^{16}$ GeV, as suggested by the running of the gauge couplings in the MSSM, naturally explains light but massive neutrinos.

Over the last several years, driven by the new and more precise neutrino data, the search for consistent SO(10) GUT models has intensified [1]. Two models, which use only small Higgs representations, are particularly interesting, namely, those by Albright and Barr (AB) [2] and by Babu, Pati and Wilczek (BPW) [3], with lopsided and hierarchical mass matrices, respectively. This choice of Higgs representations guarantees that the theory is perturbative up to the Planck scale. Then higher-dimensional operators are included to generate viable fermion masses and mixings.¹ These operators naturally produce neutrino masses at the observed scale of 0.1 eV.

The Higgs multiplets used in both models are 10_H , 16_H , $\bar{16}_H$ and 45_H . SO(10) is broken by the vacuum expectation values (vevs) $\langle 45_H \rangle \propto (B - L)$ and $\langle 16_H \rangle$ and $\langle \bar{16}_H \rangle$ in the SU(5) singlet direction. Then the SU(2)_L-doublets in 10_H and 16_H^d acquire electroweak vevs. (Since the AB model contains two distinct fields and the BPW model only one, we use 16_H^d to denote the electroweak vev of the SU(2)_L-doublet component of 16_H and $16_H'$ in the BPW and AB models, respectively.) With only $16_i 16_j 10_H$, the fermion mass matrices would coincide, $h^u = h^d = h^e = h^\nu$, and the CKM matrix would be unity. The dimension-five operator $\mathcal{O}^{B-L} \equiv 16_i 16_j 10_H 45_H$ couples differently to quarks and leptons, yielding $h^u = h^d \neq h^e = h^\nu$. Next, $\mathcal{O}^d \equiv 16_i 16_j 16_H 16_H^d$ contributes to down quarks and charged leptons only and enables a non-trivial CKM matrix. Finally, Majorana neutrino masses are gen-

erated via couplings $\mathcal{O}^N \equiv 16_i 16_j \bar{16}_H \bar{16}_H$.

Both models agree very well with observation and have been studied with increasing sophistication [5]. Thus one might ask if these models are unique or if there are more general classes of models that naturally yield the same fermion masses and mixings. In this letter, we consider all dimension-five operators that can appear with the present fields, namely $\mathcal{O}^N$, $\mathcal{O}^{B-L}$, $\mathcal{O}^d$ and $\mathcal{O}^u \equiv 16_i 16_j \bar{16}_H \bar{16}_H^u$ so that the couplings for the Dirac mass matrices read

$$h_{ij} 16_i 16_j 10_H + \frac{1}{M} \left[a_{ij} 16_i 16_j 10_H 45_H + g_{ij} 16_i 16_j 16_H 16_H^d + g'_{ij} 16_i 16_j \bar{16}_H \bar{16}_H^u \right]. \quad (1)$$

Similarly to 16_H^d , $\bar{16}_H^u$ can acquire an electroweak vev and then $\mathcal{O}^u$ contributes to up quarks and neutrinos. Whether one or both vevs of 16_H^d and $\bar{16}_H^u$ are different from zero, depends on the particular superpotential. In the AB model, a non-zero vev for $\bar{16}_H^u$ is not allowed by the F-flat direction constraints on the Higgs superpotential. In the BPW model, $\bar{16}_H^u$ can, in general, have a non-vanishing vev; however, $\mathcal{O}^u$ is suppressed due to the different charges of 16_H^d and $\bar{16}_H^u$ under the U(1) family symmetry. In this letter, we assume that with a different choice of family symmetries, both vevs are non-zero and $\mathcal{O}^u$ is present. We use the mass matrices of the AB and BPW models to study the effects of a non-vanishing contribution from the operator $\mathcal{O}^u$ on fermion masses and mixings. For simplicity, we restrict ourselves to just the second and third generations.

Hierarchical Models. We start with the case of the BPW model, with only one set of Higgs fields in the spinor representation and small contributions from the dimension-five operators. Let us assume that the contributions of $\mathcal{O}^{u,d}$ arise from integrating out heavy 10 fields

¹This is similar to SU(5) where higher-dimensional operators are used for a consistent model [4].

$[\mathcal{O}_{10}^{u,d}]$, as is the case in the AB and BPW models; we will discuss the general case later. We denote the couplings as

$$\begin{aligned} h_{33} &= 1, & g_{23} \frac{\langle 16_H \rangle}{M} \tan \gamma^d &= \eta, & a_{23} \frac{\langle 45_H \rangle}{M} &= \epsilon, \\ h_{23} &= \sigma, & g'_{23} \frac{\langle \bar{16}_H \rangle}{M} \tan \gamma^u &= \delta_{10}, \end{aligned} \quad (2)$$

with

$$\tan \gamma^d \equiv \frac{\langle 16_H^d \rangle}{\langle 10_H^d \rangle}, \quad \tan \gamma^u \equiv \frac{\langle \bar{16}_H^u \rangle}{\langle 10_H^u \rangle}; \quad (3)$$

the (22)-couplings are assumed to be sufficiently small so that they can be ignored. In the BPW model, $\delta_{10} \equiv 0$ and $\sigma, \epsilon, \eta \sim 0.1$.

The mass matrices are given by

$$\begin{aligned} h^u &= \begin{pmatrix} 0 & \epsilon + \sigma + \delta^u \\ -\epsilon + \sigma + \delta^u & 1 \end{pmatrix}, \\ h^d &= \begin{pmatrix} 0 & \epsilon + \sigma + \eta \\ -\epsilon + \sigma + \eta & 1 \end{pmatrix}, \\ h^\nu &= \begin{pmatrix} 0 & -3\epsilon + \sigma + \delta^\nu \\ 3\epsilon + \sigma + \delta^\nu & 1 \end{pmatrix}, \\ h^e &= \begin{pmatrix} 0 & -3\epsilon + \sigma + \eta \\ 3\epsilon + \sigma + \eta & 1 \end{pmatrix}, \end{aligned} \quad (4)$$

where $\delta^u = \delta^\nu = \delta_{10}$. Eqs. (4) are defined in the basis where the $\text{SU}(2)_L$ -doublets are on the left and the singlets on the right. The matrices coincide with those of the BPW model with (see Eq. (15) in Ref. [3])

$$\epsilon_{\text{BPW}} = \epsilon, \quad \sigma_{\text{BPW}} = \sigma + \delta^u, \quad \eta_{\text{BPW}} = \sigma + \eta. \quad (5)$$

Thus any realization within this class of models can reproduce the fermion mass matrices of the BPW model. In principle, one could set any one of σ, η (i.e. the operator $\mathcal{O}^d$) or δ^u (i.e. $\mathcal{O}^u$) to zero.

In Eqs. (4), we assumed that $\mathcal{O}^{u,d}$ was obtained using heavy 10 fields. Unlike $\mathcal{O}^d$, the operator $\mathcal{O}^u$ can arise from integrating out heavy singlets or fields in the adjoint representation as well. Integrating out heavy singlet fields $[\mathcal{O}_1^u]$ gives only terms to h^ν , and $\mathcal{O}_{45}^u$ contributes to h^u and h^ν in the ratio 8 : 3 [6]. Since $\delta^u \neq \delta^\nu$, we expect this more general class of models to differ from the BPW model but the differences appear in h^ν and affect the neutrino masses and mixings only. The atmospheric mixing angle is given by

$$\tan \theta_{\mu\tau}^\nu = \sqrt{\left| \frac{m_{\nu_2}}{m_{\nu_3}} \right|} \simeq \frac{h_{23}^N (3\epsilon - \sigma - \delta^\nu)}{h_{33}^N (3\epsilon + \sigma + \delta^\nu) - 2h_{23}^N}, \quad (6)$$

²In order to have a consistent notation, we replace $\epsilon/3$ and σ in Refs. [2] by $-\epsilon$ and η ; furthermore the mass matrices in Refs. [2] are defined in the basis with the singlets on the left and the doublets on the right.

where $h_{23}^N \sim \delta_1 + \delta_{45}$ and $\frac{h_{23}^N}{h_{33}^N} \equiv y_{\text{BPW}}$ (see Eq. (22) in Ref. [3]). In the BPW model, $\delta^\nu = \delta^u$ and $h_{33}^N = 1$.

Finally, let us remark that for two distinct fields, $\bar{16}_H$ and $\bar{16}_H^u$, the contribution to h^ν is not symmetric, since the neutrino singlet couples to $\bar{16}_H$ and the lepton doublet to $\bar{16}_H^u$. Additionally, if we allow heavy 120 fields, the contributions to $\mathcal{O}_{120}^{u,d}$ are not symmetric either.

Lopsided Models. Let us turn to models where the (23) and (32)-elements of the mass matrices are not of the same order. In the AB model with two distinct pairs of Higgs fields in the spinor representation, the $\text{U}(1) \times \mathbb{Z}_2 \times \mathbb{Z}_2$ family symmetry allows only $[16_2 16_H^d]_{10} [16_3 16_H]_{10}$. In addition, $\sigma \equiv 0$ (see Eqs. (4)). As a result, h^d and h^e have lopsided structures, and a large value for $\tan \gamma^d$ leads to $\eta \simeq 1.8$, whereas $\epsilon \simeq 0.05$ [2].

Let us assume that a specific family symmetry leads to h^d and h^e as in the AB model and that $\mathcal{O}^u$ contributes to h^u and h^ν . Then the quark matrices read²

$$h^u = \begin{pmatrix} 0 & \epsilon + \delta^u \\ -\epsilon + \delta^u & 1 \end{pmatrix}, \quad h^d = \begin{pmatrix} 0 & \epsilon \\ -\epsilon + \eta & 1 \end{pmatrix}, \quad (7)$$

with $\delta^u = \delta_{10} + \delta_{45}$. In the AB model, $\delta^u \equiv 0$. Note that the contribution from $\mathcal{O}^u$ to h^u is symmetric, regardless of the family symmetry, because the quark doublet and the up-quark singlet belong to the same $\text{SU}(5)$ -representation. In contrast, the down quark and lepton doublets are in different $\text{SU}(5)$ -representations than the singlets.

To obtain a viable hierarchy for the up quarks, δ^u must be small. Then we obtain

$$\frac{m_c}{m_t} \simeq |\delta^2 - \epsilon^2|, \quad \frac{m_s}{m_b} \simeq \frac{\epsilon\eta}{1 + \eta^2}, \quad V_{cb} \simeq \frac{\epsilon\eta^2}{1 + \eta^2} + \delta. \quad (8)$$

Unlike in the BPW case, we have found an additional parameter in h^u , which cannot be absorbed in the original parameters and needs to be included in the fit. On the other hand, this parameter has to be small in order to reproduce the quark masses and mixing. We do not expect significant changes to the neutrino sector, in particular the mixing angle is still dominated by the lopsidedness of h^e .

For the hierarchical models, we found that we could set $\eta = 0$ and still reproduce the results of the BPW model with $\delta^u \equiv 0$ (cf. Eq. (5)). Thus one might consider the case $\eta = 0$ for the AB-type models with mass matrices as

displayed in Eqs. (7); however, they rely on the lopsidedness of h^d and h^e . Hence, one could try to mimic these effects with a lopsided matrix h^u .

A lopsided structure for h^u can be generated if we take into account antisymmetric contributions and tune them such that different terms in the mass matrices cancel. We already know that $\mathcal{O}^{B-L}$ gives antisymmetric contributions (see Eqs. (4)) but $\epsilon \sim 0.1$ is too small to generate lopsided entries. As already mentioned, we can also integrate out heavy fields in the (large) 120 representation, and $\mathcal{O}_{120}^u = [16_i 16_j]_{120} [\overline{16}_H \overline{16}_H^u]_{120}$ yields an antisymmetric contribution, δ_{120} . If $\delta_{120} = -\delta$, then

$$h^u = \begin{pmatrix} 0 & \epsilon \\ -\epsilon + 2\delta & 1 \end{pmatrix}, \quad (9)$$

and we get

$$\frac{m_c}{m_t} \simeq \frac{2\epsilon\delta}{1+4\delta^2}, \quad V_{cb} \simeq \epsilon \left(\frac{1}{1+4\delta^2} - \frac{1}{1+\eta^2} \right). \quad (10)$$

Compared to Eqs. (7), the factor $\frac{2\delta}{1+4\delta^2}$ must be of order ϵ in order to enable the up quark mass hierarchy. The two solutions are $\delta \sim \epsilon$ and $\delta \sim \frac{1}{\epsilon}$, but if $\eta = 0$, we obtain $V_{cb} \simeq \frac{4\epsilon\delta^2}{1+4\delta^2}$ so that $\delta \simeq 1$. Hence, $\eta = 0$ is not a viable case.

With both δ and η non-zero, one should consider what the natural scale for δ is. Since $\frac{\langle 10_H^u \rangle}{\langle 10_H^d \rangle}$ can be much bigger than one, a large value for $\tan \gamma^d$ is not unnatural [2]. On the other hand, we expect $\tan \gamma^u$ not to be bigger than one so that $\delta \sim \epsilon \sim 0.1$ is a natural choice. Therefore we do not consider solutions with $\delta \sim \frac{1}{\epsilon} \gg 1$.

Conclusions. We have shown that the BPW model belongs to a class of models with equivalent fermion masses and mixings; it represents the special case where the operator $\mathcal{O}^u$ vanishes. If we factor in heavy singlets and heavy fields in the adjoint representation, then $\mathcal{O}^u$ does not contribute equally to h_u and h_ν . Therefore we expect deviations from the BPW model in the neutrino sector, which

are presumably small. In AB-type models, including $\mathcal{O}^u$ generically introduces a new parameter in the quark mass matrices which is smaller than the contribution from the operator $\mathcal{O}^d$. Unlike the down quarks and leptons, family symmetries cannot generate a lopsided up-quark mass matrix unless we allow for fine-tuning.

We restricted the discussion to the second and third generations only; however, the results apply to the three-generational case as well.

We are grateful for valuable discussions with S. Wiltenbrock and for useful comments on the manuscript by C. Albright and J. Pati. This work was supported in part by the U. S. Department of Energy under contract No. DE-FG02-91ER40677.

References

- [1] For recent reviews on SO(10) models, see J. C. Pati, hep-ph/0507307; A. Melfo and G. Senjanovic, hep-ph/0511011.
- [2] C. H. Albright and S. M. Barr, Phys. Rev. D **58**, 013002 (1998); C. H. Albright, K. S. Babu and S. M. Barr, Phys. Rev. Lett. **81**, 1167 (1998); C. H. Albright and S. M. Barr, Phys. Lett. B **452**, 287 (1999); *ibid* **461**, 218 (1999); Phys. Rev. Lett. **85**, 244 (2000); Phys. Rev. D **62**, 093008 (2000).
- [3] K. S. Babu, J. C. Pati and F. Wilczek, Nucl. Phys. B **566**, 33 (2000).
- [4] S. Wiesenfeldt, Mod. Phys. Lett. A **19**, 2155 (2004) and references therein.
- [5] C. H. Albright and S. M. Barr, Phys. Rev. D **64**, 073010 (2001); *ibid* **69**, 073010 (2004); C. H. Albright, Phys. Rev. D **72**, 013001 (2005); K. S. Babu, J. C. Pati and P. Rastogi, Phys. Rev. D **71**, 015005 (2005); Phys. Lett. B **621**, 160 (2005); P. Rastogi, Phys. Rev. D **72**, 075002 (2005).
- [6] K. S. Babu and S. M. Barr, Phys. Rev. Lett. **85**, 1170 (2000); P. Nath and R. M. Syed, Nucl. Phys. B **618**, 138 (2001).